

MONITORING AEROSOL PRESENCE OVER A 15-YEAR PERIOD USING THE ABSORBING AEROSOL INDEX MEASURED BY GOME-1, SCIAMACHY, AND GOME-2

L. G. Tilstra¹, M. de Graaf^{1,2}, O. N. E. Tuinder¹, R. J. van der A¹, and P. Stammes¹

¹Royal Netherlands Meteorological Institute (KNMI), P.O. Box 201, 3730 AE De Bilt, The Netherlands

²Delft University of Technology, P.O. Box 5, 2600 AA Delft, The Netherlands

Email: tilstra@knmi.nl, graafdem@knmi.nl, tuinder@knmi.nl, avander@knmi.nl, stammes@knmi.nl

ABSTRACT

In this paper we study 15-year long time series of the Absorbing Aerosol Index (AAI) for a selection of the most prominent aerosol producing regions on the globe. The time series of the AAI were recorded by the satellite instruments GOME-1, SCIAMACHY, and GOME-2 aboard the ERS-2, Envisat, and MetOp-A satellites, respectively. These three satellite instruments suffer, each in their own way, from severe instrument degradation in the ultraviolet (UV) wavelength range from which the AAI is derived. To be able to perform a reliable analysis on the AAI time series, we first remove the effects of instrument degradation from the Earth reflectances before calculating the AAI from them. To validate the resulting time series, we use tropospheric NO₂ data as a reference in the regions dominated by biomass burning events. From this it is found that the regional AAI data follow the regional tropospheric NO₂ data well. Therefore, it is possible to accurately remove the effects of instrument degradation and to combine the AAI data from the three satellite instruments. The time series of the AAI for the studied aerosol regions show no clear trend over the studied 15-year time period from 1995 to 2011.

Key words: aerosols; Absorbing Aerosol Index; trends.

1. ABSORBING AEROSOL INDEX

The Absorbing Aerosol Index (AAI) is an index based on a comparison of measured UV reflectances with simulated Rayleigh reflectances [1, 2]. These simulated reflectances are calculated for cloud-free and aerosol-free atmospheres in which only Rayleigh scattering, absorption by molecules, Lambertian surface reflection and surface absorption can take place. The AAI is derived from another quantity, called the residue, which is defined as

$$r = -100 \cdot 10 \log \left(\frac{R_{\lambda}^{\text{obs}}}{R_{\lambda}^{\text{Ray}}} \right). \quad (1)$$

In this equation, R^{obs} refers to reflectances measured by one of the three satellite instruments, while R^{Ray} refers to modelled Rayleigh reflectances. The symbol λ refers to the first, shortest wavelength of the AAI wavelength pair, which is 340 nm. The surface albedo A_s used in the simulations for this wavelength is assumed to be the same as the surface albedo at the second wavelength $\lambda_0 = 380$ nm. The surface albedo at 380 nm in turn is found from requiring that the simulated Rayleigh reflectance equals the measured reflectance at this wavelength. That is, we have the following two constraints:

$$R_{\lambda_0}^{\text{obs}} = R_{\lambda_0}^{\text{Ray}}(A_s) \quad ; \quad A_s(\lambda) = A_s(\lambda_0) \quad (2)$$

The two equations above basically define the algorithm that is used to calculate the residue. When a positive residue is found, absorbing aerosols were detected. Negative or zero residues on the other hand suggest an absence of absorbing aerosols. Therefore, the AAI is defined as equal to the residue where the residue is positive, and it is simply not defined for negative values of the residue.

The GOME-1, SCIAMACHY, and GOME-2 AAI are developed and produced at the KNMI and the data are available for download on the TEMIS website via the URL <http://www.temis.nl/>. The GOME-2 AAI is an operational product in the O3M SAF. For the main characteristics of the AAI products, please consult Tab. 1.

2. INSTRUMENT DEGRADATION

Instrument degradation has been shown to have a large impact on the AAI [3, 4, 5]. In the current paper we will be presenting and analysing AAI data that were corrected for instrument degradation in the manner described in the paper by Tilstra et al. [4]. In this paper, the Earth reflectance measured by SCIAMACHY is corrected for the effects of instrument degradation by analysing time series of the global mean reflectance, extracting a polynomial function that describes the effects of instrument degradation, and performing the resulting correction. In this approach, the time series of the global mean reflectance

Table 1. Main characteristics of GOME-1, SCIAMACHY, and GOME-2, of their derived AAI products, and of the time periods covered by the three AAI datasets. The AAI wavelength pair is 340/380 nm in all cases.

	Platform / Satellite	Equator passing time (local time)	Pixel size (km ²)	Days needed for global coverage	AAI wavelength pair (nm)	AAI level-2 version	AAI time period covered
GOME-1	ERS-2	10:30 a.m.	320 × 40	3	340 / 380	5.0	1995–2003
SCIAMACHY	Envisat	10:00 a.m.	60 × 30	6	340 / 380	5.0	2002–2011
GOME-2	MetOp-A	09:30 a.m.	80 × 40	1.5	340 / 380	1.18	2007–2011

are fitted with a Fourier series multiplied with a polynomial base. The Fourier terms are meant to describe the seasonal variation of the global mean reflectance. These seasonal variations are not caused by instrument degradation, and therefore disregarded. The polynomial terms, on the other hand, represent the drop in signal due to instrument degradation. Knowledge of the polynomial base allows a straightforward correction of the reflectances. The method described here was applied individually to the GOME-1, SCIAMACHY, and GOME-2 instruments. The reflectance measurements performed by SCIAMACHY were additionally corrected for radiometric calibration problems that were reported earlier [6, 7] using an approach introduced by Tilstra et al. [8].

3. STUDYING TIME SERIES OF THE AAI

In this section we will analyse time series of the AAI determined from GOME-1, SCIAMACHY, and GOME-2 observations for a selection of known aerosol regions. The box-shaped regions are defined graphically in Fig. 1.




Figure 1. Graphical representation of the aerosol regions studied: Borneo, Amazonia, the Sahel, and West Africa. Also plotted (in brown) is the averaged AAI field measured by SCIAMACHY over the years 2002–2010.

3.1. Case A: Borneo

In Fig. 2 we present the time series of the AAI for the region containing the island of Borneo as illustrated in Fig. 1. On the vertical axis we plotted the regional mean

AAI, averaged with a running mean of 31 days. The red data points were measured by GOME-1, the brown data points by SCIAMACHY, and the blue data points by GOME-2. Notice the two strong peaks in the time period 1997/1998. These were caused by forest fires due to extreme droughts in this El Niño episode. On Borneo, a total area of 120.000 km² forest was destroyed [9].

From Fig. 2 it is found that peaks occur once or twice a year: either in the months March/April or in the months September/October. The probability that such a peak occurs turns out to be higher in the more recent years. In a recent paper by Van der Werf et al. [10], time series of fire emission estimates are given for a somewhat similar region as the one indicated in Fig. 1. The agreement between the time series of fire emissions given there and the AAI shown in Fig. 2 is quite good. The baseline (background level) of the AAI time series, consisting of low values of ~ 0.2 , is rather stable. This is a good indication that instrument degradation was removed successfully. Also notice that the three data sets overlap consistently with each other where they overlap in time.

3.2. Case B: Amazonia

The mean AAI for the Amazonia region is presented in Fig. 3. There is a strong seasonal cycle, and a strong year-to-year variation. In most years, there is a strong biomass burning peak present in the months August/September. For a qualitative validation we can again make use of results from the paper by Van der Werf et al. [10]. The agreement is again quite satisfactory. To get to a more quantitative validation, we resort to AERONET data of station Alta Floresta. This station is represented by the blue dot in Fig. 1. We used level 2.0 data, and only accepted measurements which had a measurement time close to the local measurement time of GOME-1, SCIAMACHY, and GOME-2. Furthermore, we applied a running mean of 11 days. The resulting time series is presented in Fig. 4.

From a qualitative point of view there is certainly agreement, but there is also quite a poor correlation between the peak heights found in Fig. 3 and those found in Fig. 4. Efforts to improve the correlation by reducing the size of the box shown in Fig. 1, in order to make it agree more with the observational area of the AERONET station, were only partly successful. Possible explanations for




Figure 2. Regional mean AAI for the box containing the island Borneo as a function of time over the period 1995–2011.




Figure 3. Regional mean AAI for the Amazonian box as a function of time over the period 1995–2011.




Figure 4. Aerosol Optical Thickness for AERONET station “Alta Floresta”. A running mean of 11 days was applied.




Figure 5. Mean tropospheric NO_2 column for the Amazonian box as a function of time over the period 1995–2011.




Figure 6. Regional mean AAI for the Sahel region as a function of time over the period 1995–2011.




Figure 7. Mean tropospheric NO₂ column for the Sahel region as a function of time over the period 1995–2011.




Figure 8. Regional mean AAI for the box near West Africa as a function of time over the period 1995–2011.




Figure 9. Mean tropospheric NO₂ column for the box near West Africa as a function of time over the period 1995–2011.

the remaining differences are the cloud screening in the AERONET level 2.0 data and the fact that the AAI and AOT are by definition not related by a fixed unique relationship. Also the large amount of missing AERONET data might play a role in the discrepancies found.

In a next attempt to validate the AAI time series we analyse tropospheric NO₂ data measured by the instruments GOME-1 and SCIAMACHY for the same region and for the same time period. Emission of NO₂ by biomass burning processes is well-known [11, 12]. Since the area studied is a typical biomass burning region, we would expect some sort of correlation between the concentration of aerosols and that of the tropospheric NO₂ column (as found in Veefkind et al. [13]). In Fig. 5 we present the tropospheric NO₂ column, measured by GOME-1 and SCIAMACHY, averaged over the studied region and treated with a running mean of 31 days. The GOME-1 data are plotted using the light blue colour and the dark blue colour is used to denote the SCIAMACHY data.

The correlation between the AAI time series presented in Fig. 3 and the tropospheric NO₂ column shown in Fig. 5 is quite good, even on a quantitative level. This indicates that the seasonal variations in the aerosol presence are well captured by the AAI, but it also yields confidence in the recorded year-to-year variations in these.

3.3. Case C: The Sahel

The aerosol presence in the Sahel region is known to be driven by biomass burning events and/or dust storms, depending on the time of the year [14, 15]. The time series of the regional mean AAI is given in Fig. 6. The aerosol presence has a peak in the period between December and March. Looking at the time series of the tropospheric NO₂ column presented in Fig. 7, we conclude that this period is dominated by biomass burning. The agreement between the AAI and the tropospheric NO₂ column is again good. The time series of the AAI again shows a stable baseline, supporting the idea that no remnants of instrument degradation have influenced the time series.

The time series of the three instruments again overlap nicely in the overlap regions. A clear trend in the time series of the regional mean AAI cannot be found.

3.4. Case D: West Africa

The aerosol presence in West Africa is well known to be dominated by biomass burning events for most of the year. The time series of the regional mean AAI is presented in Fig. 8. The seasonal variation is quite strong, showing a maximum in the period between July and September, and a secondary local maximum in the month of January. This secondary maximum is presumably related to transport of desert dust from the Sahel region. The year-to-year variation in Fig. 8 is quite modest.

The time series of the mean tropospheric NO₂ column for the same region is shown in Fig. 9. The correlation between this time series and the one presented in Fig. 8 is very high. The shapes of the seasonal variation are quite similar. The only difference is that the secondary maxima in the AAI time series do not exist in the NO₂ time series. This would support the idea that the secondary maxima in the AAI time series are indeed not related to biomass burning aerosols, but to desert dust aerosols.

Next, in Fig. 10, we present the scatter plot of the regional mean residue versus the regional mean tropospheric NO₂ column. The correlation is surprisingly good, which we attribute to the averaging over space and time. The averaging over space and time is needed to compensate for the different atmospheric trajectories followed by the aerosols and the NO₂ molecules. We conclude that the seasonal variation of the AAI time series is very similar to the seasonal variation of the time series of the tropospheric NO₂ column. Next to this, the year-to-year variation of the AAI seems to be correct.




Figure 10. Regional mean tropospheric NO₂ column versus the regional mean residue for the box near West Africa. There is a clear correlation, at least in the residue regime where aerosols are expected to have been present.

4. CONCLUSIONS

In this paper we showed that it is possible to successfully combine Absorbing Aerosol Index (AAI) data from GOME-1, SCIAMACHY, and GOME-2 into a 15-year long data archive of aerosol presence. To be able to achieve this, instrument degradation must be accurately corrected for. Using the AAI data, we studied time series of aerosol presence for four typical aerosol regions. The time series showed no apparent trend, indicating the correctness of the applied correction for instrument degradation. In the overlap periods of the different satellite instruments there was good agreement. The time series were compared with, amongst other things, time series of

tropospheric NO₂ data. For scenes controlled by biomass burning events, the agreement was quite good. From this we conclude that we are indeed able to correct for instrument degradation in a sufficient manner. This allows accurate long-term studies of aerosol presence using the AAI. We also anticipate that the combination of the tropospheric NO₂ column with the AAI can be used to verify chemical-transport-models, along the same lines as was done by Veefkind et al. [13].

ACKNOWLEDGEMENTS

The work presented in this paper was financed by EUMETSAT via the O3M SAF, and by the Netherlands Space Office (NSO) through the SCIA-Visie project. We acknowledge the use of AERONET data and thank Brent Holben for his effort in establishing and maintaining the Alta Floresta site. We also acknowledge the free use of tropospheric NO₂ column data from the GOME and SCIAMACHY sensors from www.temis.nl.

REFERENCES

- [1] Herman, J.R., Bhartia, P.K., Torres, O., Hsu, C., Seftor, C. & Celarier, E. (1997). Global distributions of UV-absorbing aerosols from Nimbus 7/TOMS data. *J. Geophys. Res.* **102**(D14), 16,911–16,922, doi:10.1029/96JD03680.
- [2] Torres, O., Bhartia, P.K., Herman, J.R., Ahmad, Z. & Gleason, J. (1998). Derivation of aerosol properties from satellite measurements of backscattered ultraviolet radiation: Theoretical basis. *J. Geophys. Res.* **103**(D14), 17,099–17,110, doi:10.1029/98JD00900.
- [3] Tilstra, L.G., Tuinder, O.N.E. & Stammes, P. (2010). GOME-2 Absorbing Aerosol Index: statistical analysis, comparison to GOME-1 and impact of instrument degradation. In *Proc. 2010 'EUMETSAT Meteorological Satellite Conference'*, EUMETSAT P.57, EUMETSAT, Darmstadt, Germany.
- [4] Tilstra, L.G., De Graaf, M., Aben, I. & Stammes, P. (2012). In-flight degradation correction of SCIAMACHY UV reflectances and Absorbing Aerosol Index. *J. Geophys. Res.* **117**, D06209, doi:10.1029/2011JD016957.
- [5] Tilstra, L.G., Tuinder, O.N.E. & Stammes, P. (2012). A new method for in-flight degradation correction of GOME-2 Earth reflectance measurements, with application to the Absorbing Aerosol Index. In *Proc. 2012 'EUMETSAT Meteorological Satellite Conference'*, EUMETSAT P.61, EUMETSAT, Darmstadt, Germany.
- [6] Tilstra, L.G. & Stammes, P. (2006). Intercomparison of reflectances observed by GOME and SCIAMACHY in the visible wavelength range. *Appl. Opt.* **45**(17), 4129–4135, doi:10.1364/AO.45.004129.
- [7] Tilstra, L.G. & Stammes, P. (2007). Earth reflectance and polarization intercomparison between SCIAMACHY onboard Envisat and POLDER onboard ADEOS-2. *J. Geophys. Res.* **112**, D11304, doi:10.1029/2006JD007713.
- [8] Tilstra, L.G., Van Soest, G. & Stammes, P. (2005). Method for in-flight satellite calibration in the ultraviolet using radiative transfer calculations, with application to Scanning Imaging Absorption Spectrometer for Atmospheric Cartography (SCIAMACHY). *J. Geophys. Res.* **110**, D18311, doi:10.1029/2005JD005853.
- [9] Wooster, M.J. & Strub, N. (2002). Study of the 1997 Borneo fires: Quantitative analysis using global area coverage (GAC) satellite data. *Global Biogeochem. Cycles* **16**(1), 9-1–9-12, doi:10.1029/2000GB001357.
- [10] Van der Werf, G.R., Randerson, J.T., Giglio, L., Collatz, G.J., Mu, M., Kasibhatla, P.S., Morton, D.C., DeFries, R.S., Jin, Y. & Van Leeuwen, T.T. (2010). Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmos. Chem. Phys.* **10**, 11707–11735, doi:10.5194/acp-10-11707-2010.
- [11] Van der A, R.J., Peters, D.H.M.U., Eskes, H., Boersma, K.F., Van Roozendael, M., De Smedt, I. & Kelder, H.M. (2006). Detection of the trend and seasonal variation in tropospheric NO₂ over China. *J. Geophys. Res.* **111**, D12317, doi:10.1029/2005JD006594.
- [12] Van der A, R.J., Eskes, H.J., Boersma, K.F., Van Noije, T.P.C., Van Roozendael, M., De Smedt, I., Peters, D.H.M.U. & Meijer, E.W. (2008). Trends, seasonal variability and dominant NO_x source derived from a ten year record of NO₂ measured from space. *J. Geophys. Res.* **113**, D04302, doi:10.1029/2007JD009021.
- [13] Veefkind, J.P., Boersma, K.F., Wang, J., Kurosu, T.P., Krotkov, N., Chance, K. & Levelt, P.F. (2011). Global satellite analysis of the relation between aerosols and short-lived trace gases. *Atmos. Chem. Phys.* **11**, 1255–1267, doi:10.5194/acp-11-1255-2011.
- [14] Tilstra, L.G., De Graaf, M., Aben, I. & Stammes, P. (2007). Analysis of 5 years SCIAMACHY Absorbing Aerosol Index data. In *Proc. 2007 'Envisat Symposium'* (Eds. H. Lacoste & L. Ouweland), ESA SP-636 (CD-ROM), ESA Communication Production Office, European Space Agency, Noordwijk, The Netherlands.
- [15] De Graaf, M., Tilstra, L.G., Aben, I. & Stammes, P. (2010). Satellite observations of the seasonal cycles of absorbing aerosols in Africa related to the monsoon rainfall, 1995–2008. *Atmos. Environ.* **44**(10), 1274–1283, doi:10.1016/j.atmosenv.2009.12.038.