

Geophysical Research Letters



RESEARCH LETTER

10.1029/2020GL089912

Special Section:

The COVID-19 Pandemic:
Linking Health, Society and
Environment

Key Points:

- NO_x emissions derived from TROPOMI observations show reductions for individual Chinese cities of about 35% due to the COVID-19 lockdown
- Emissions of coal power plants and maritime transport show strong reductions (25–40%) during the lockdown
- Urban emissions rebound in March to levels before the lockdown, while emissions of power plants and maritime transport take longer to recover

Supporting Information:

- Supporting Information S1

Correspondence to:

J. Ding,
jieding.ding@knmi.nl

Citation:

Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrakou, T., van Geffen, J. H. G. M., et al. (2020). NO_x emissions reduction and rebound in China due to the COVID-19 crisis. *Geophysical Research Letters*, 46, e2020GL089912. <https://doi.org/10.1029/2020GL089912>

Received 21 JUL 2020

Accepted 29 AUG 2020

Accepted article online 9 SEP 2020

NO_x Emissions Reduction and Rebound in China Due to the COVID-19 Crisis

J. Ding¹ , R. J. van der A^{1,2} , H. J. Eskes¹ , B. Mijling¹ , T. Stavrakou³ ,
J. H. G. M. van Geffen¹ , and J. P. Veefkind^{1,4}

¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands, ²School of Atmospheric Physics, Nanjing University of Information Science and Technology (NUIST), Nanjing, China, ³Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium, ⁴Department of Geoscience and Remote Sensing, Delft University of Technology, Delft, The Netherlands

Abstract During the COVID-19 lockdown (24 January–20 March) in China low air pollution levels were reported in the media as a consequence of reduced economic and social activities. Quantification of the pollution reduction is not straightforward due to effects of transport, meteorology, and chemistry. We have analyzed the NO_x emission reductions calculated with an inverse algorithm applied to daily NO₂ observations from TROPOMI onboard the Copernicus Sentinel-5P satellite. This method allows the quantification of emission reductions per city and the analysis of emissions of maritime transport and of the energy sector separately. The reductions we found are 20–50% for cities, about 40% for power plants, and 15–40% for maritime transport depending on the region. The reduction in both emissions and concentrations shows a similar timeline consisting of a sharp reduction (34–50%) around the Spring festival and a slow recovery from mid-February to mid-March.

Plain Language Summary During the COVID-19 lockdown in China, air quality had strongly improved. Here we study what sources were reduced and how much the reduction per city was. We used TROPOMI observations of the Sentinel-5P satellite, which monitors the Earth's atmosphere daily. We focused on observations of the pollutant “nitrogen dioxide,” an important precursor of air pollution in the atmosphere. With our novel methodology we are able to calculate the pollution back to the sources of the emissions, whether these are big cities, industrial regions, power plants, or busy shipping lanes. We applied this method to East China, where the 36 biggest Chinese cities are located. Almost all those cities showed strong emission reductions of 20–50% during the lockdown in February 2020. Besides urban China, we found an average emission reduction of 40% over coal power plants and a reduction in maritime transport by 15–40% depending on the region. The period of reduced emissions lasted until around the end of February, and the emissions slowly returned to normal during the month March 2020. Exception is the region Wuhan, the center of the COVID-19 crisis, where emissions started to rebound since 8 April, the end of their lockdown period.

1. Introduction

The year 2020 is an unprecedented year, with the novel coronavirus, causing the COVID-19 disease spreading over the whole world, infecting millions of people and causing hundreds of thousands of fatalities (WHO, 2020). On 11 March 2020, the World Health Organization (WHO) qualified the spread of COVID-19 as a pandemic. To prevent the spread of the disease, many affected countries implemented COVID-19 regulations. China, the first country facing the outbreak of COVID-19, enacted a lockdown from 24 January to 20 March 2020 in the Hubei province where the first cases were reported from its capital Wuhan, while other provinces limited all outdoor activities since the Chinese New Year and gradually resumed the work after 10 February (Tian et al., 2020; Wang et al., 2020).

The strict COVID-19 regulations lead to a reduction of road and air traffic, a temporary closing of companies, and a decrease of industrial productivity. These in consequence affect emissions of air pollutants, especially from the transport and industry sectors, which are significant sources of NO_x (NO_x = NO₂ + NO) in cities. Several studies presented a large decrease of NO₂ concentration during the lockdown period in China from both in situ and satellite observations (Huang et al., 2020; Wang et al., 2020). Tropospheric NO₂ column concentrations observed by the TROPOMI (TROPOspheric Monitoring Instrument) on the Sentinel-5P satellite

©2020. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure 1. TROPOMI NO₂ columns over East China after the Chinese New Year in 2019 (a) and 2020 (b). NO_x emissions for the same period in 2019 (c) and 2020 (d) derived with DECSO.

decrease about 35% over China and some areas up to 60% during the COVID-19 regulation period compared to the same period of 2019 (Bauwens et al., 2020; Liu et al., 2020; Zhang et al., 2020). In March 2020, after the resumption of work and the gradual lifting of the lockdown restrictions, the NO₂ concentrations quickly increased to similar levels as in the previous year (Bauwens et al., 2020). Because NO₂ concentrations are

affected by meteorology, chemistry, and transport, large concentration variations are expected from day to day. Therefore, the concentrations alone provide only an indication of the impact of the COVID-19 measures on air pollution. Bottom-up inventories are usually updated with few years delay due to the complexity of gathering all statistic information on source sector, land-use, and sector-specific emission factors. A top-down approach using satellite observations has been demonstrated to be able to accurately and quickly provide emission estimates (Miyazaki et al., 2020; Stavrou et al., 2013). Here we derived the NO_x emissions by using the satellite observations and a chemistry-transport model (CTM). The model is driven by meteorological analyses, accounting for the weather-related variability. The high spatial resolution of the TROPOMI observations and the inverse modeling system allows us to quantify the impact of the COVID-19 measures and distinguish emissions from cities, power plants, and maritime transport separately. Recently, NO_x emissions derived from the high resolution NO_2 observations of TROPOMI have been reported by Goldberg et al. (2019) and van der A et al. (2020).

To this purpose, we use the Daily Emission estimates Constrained by Satellite Observations (DECISO) algorithm, which has been demonstrated to capture emission changes in a short time period at city level (Ding et al., 2015; Mijling & van der A, 2012). This study presents NO_x emissions estimated from Sentinel-5P TROPOMI observations from 2019 to April 2020 over East Asia. The high spatial resolution satellite observations and daily global coverage allow us to monitor fast emission changes per city due to the implementation and to the relaxing of COVID-19 regulations.

2. Methodology

2.1. NO_2 Observations by TROPOMI

The Copernicus Sentinel-5P satellite carries the TROPOMI instrument (Veefkind et al., 2012). TROPOMI is a spectrometer combining a high spectral resolution with high spatial resolution ($3.5 \times 5.5 \text{ km}^2$ at nadir for the NO_2 observations), low noise, and a daily global coverage. Despite the much smaller footprints, the spectral fits of the individual TROPOMI ground pixels have 30% smaller noise than those from the earlier Ozone Monitoring Instrument (OMI), and the average values agree well within 5% (van Geffen et al., 2020).

Validation studies (Judd et al., 2020; Tack et al., 2020; Verhoelst et al., 2020) show that the currently available TROPOMI product (versions 1.2.2 and 1.3.0) has tropospheric columns with effectively a typical systematic bias of about -15% (see Supporting Information), and we expect the derived emissions from these observations to be low by such an amount on average. In the relative comparisons discussed in this paper for both columns and emissions we expect a large part of such a multiplicative bias to cancel out.

Figures 1a and 1b show the mean TROPOMI NO_2 tropospheric column observations gridded on a 0.02° by 0.02° grid for the periods 8–28 February 2020 compared with 18 February–4 March 2019, both after the Chinese New Year holidays. Very prominent concentration reductions are observed in 2020 compared to 2019.

The TROPOMI tropospheric NO_2 columns are pre-processed into “super-observations,” representing the integrated average of the TROPOMI observations over the $0.25^\circ \times 0.25^\circ$ grid cells of the model after filtering for clouds. The basic concept of super-observations has been explained in Miyazaki et al. (2012) and Boersma et al. (2016). They have shown that clustering individual observations into super-observations has a positive impact on the analysis. The super-observation error takes into account spatial correlations between individual TROPOMI observations as well as representativity errors in the case of incomplete coverage. Averaging kernels are also computed for these super-observations and are used in the emission estimates described below. This has the advantage that the inversion result becomes independent of the coarser-resolution of the a priori profile used in the retrieval of the tropospheric column.

2.2. NO_x Emissions From DECISO

DECISO is a state-of-the-art inverse algorithm developed by Mijling and van der A (2012) to update daily emissions of short-lived atmospheric constituents using an extended Kalman filter in which emissions are translated to concentrations via a CTM and compared to the satellite observations. The sensitivity of concentrations to emissions is calculated from a trajectory analysis to account for transport of the short-lived gas by using a single CTM forward run. DECISO has been successfully applied to NO_2 observations from OMI and

TROPOMI over different regions (Ding et al., 2017, 2018; Mijling & van der A, 2012; van der A et al., 2020). In this study, daily NO_x emissions from January 2019 to April 2020 over East Asia ($102\text{--}120^\circ\text{E}$, $18\text{--}50^\circ\text{N}$) are derived with DECSO using the Eulerian regional off-line CTM CHIMERE v2013 (Menut et al., 2013) and TROPOMI NO_2 observations. The implementation of CHIMERE v2013 in DECSO is described in Ding et al. (2015). The latest development and validation of DECSO are presented in previous studies (Ding et al., 2017; van der A et al., 2020). In our current approach, we apply DECSO to the super-observations of TROPOMI instead of directly using individual TROPOMI observations. Figures 1c and 1d show the mean NO_x emissions derived from TROPOMI for the same period as Figures 1a and 1b in 2019 and 2020 after the Chinese New Year. We see lower NO_x emissions in February 2020.

2.3. In Situ Observations

More than 1,500 in situ stations covering all major cities in China are operated by the China National Environmental Monitoring Center. They provide hourly observations of the pollutants PM_{10} , $\text{PM}_{2.5}$, O_3 , NO_2 , SO_2 , and CO (Bai et al., 2020). NO_2 is measured by a chemiluminescence technique (Zhang & Cao, 2015). Data can be accessed via websites of third parties (such as <http://www.pm25.in> and <http://www.aqicn.org>). For this study we have averaged the various in situ NO_2 observations in a city to a single value per hour for each of 36 selected major cities. For comparison with model results, we calculated a daily value based on the observations from 10:00 to 18:00 local time. The daytime selection is due to large inaccuracies in simulations of the nighttime boundary layer height.

2.4. Ensemble Modeling

An operational multi-model forecasting system for air quality has been developed to provide air quality services for urban areas of China (Brasseur et al., 2019; Petersen et al., 2019). This system has been developed within the EU-funded FP-7 projects: MarcoPolo and PANDA. The ensemble model system includes nine global and regional chemistry-transport models from different research institutes from Europe and China. The ensemble service has a typical resolution of about 20 km. It provides daily forecasts of ozone, nitrogen oxides, and particulate matter for the 36 largest urban areas of East China (i.e., population higher than 3 million according to the census of 2010; NBS, 2010). These individual 3-day forecasts as well as the mean and median concentrations are publicly accessible (<http://www.marcopolo-panda.eu>). The emission inventories used as input to the models of the ensemble do not account for the Chinese New Year or the COVID-19 lock down period. Therefore, the ensemble model represents the business-as-usual scenario.

3. NO_x Emissions Reductions

NO_x emissions have been affected since the strict regulations started in China, especially in Hubei. We select three periods to quantify the impact of the COVID-19 regulations. The first period (P1) is 3 weeks before the implementation of the COVID-19 regulations, 3–23 January in 2020, which is also just before the Chinese New Year. The second period (P2) is 8–28 February, which is regarded as the regulation period. The third period (P3) is from 18 March to 7 April, when most regions in China resumed working. We calculated the average of NO_x emissions derived with DECSO in each period and compare their differences. Figure 2 shows the relative changes of NO_x emissions during the selected three periods over the grid cells with high anthropogenic (above $3 \text{ kg N/km}^2/\text{day}$) NO_x emissions. We observe a strong decrease by at least 30% of NO_x emissions over China in P2 compared to P1 (Figure S1 shows the emission changes on provincial level). A few grid cells with increased emissions often coincide with industrial areas. In P3, NO_x emissions increased compared to P2 but are still lower than in P1 because of the step-wise resumption of work and social life. The NO_x emissions in South Korea are not significantly changed in P2 compared to the changes in China during the three periods (Figure S1), because South Korea adopted less restrictive COVID-19 regulations, mostly on voluntary basis (Bauwens et al., 2020). In Figure 2, we see that the NO_x emissions over sea also decrease. We calculate the NO_x emissions over the ship lanes over Chinese seas defined in the study of Ding et al. (2018). The emissions due to sea-transport from Shanghai to Guangzhou are less affected than the transport over land and are found to decrease by about 25% in P2 and increase again with 18% in P3 in comparison to P2. A more significant emission decline was found in the Yellow Sea and Bohai area, where NO_x emissions reduced by about 41% in P2 and continued decreasing by 6% in P3.




Figure 2. The relative difference in NO_x emissions between (a) P2 and P1, (b) P3 and P2, and (c) P3 and P1. P1 is 3–23 January. P2 is 8–28 February. P3 is 18 March–7 April. The changes in emissions are shown in the figure for emissions higher than 3 kg(N)/km²/day in P1 to remove areas with dominating biogenic emissions or rural areas.

At city level changes in NO_x emissions started from January 2019. Figure 3 shows the time series of emissions at six large cities in China and in Seoul, the capital of South Korea. We infer a very strong NO_x emission decrease of more than 50% during and after the 2020 Chinese New Year in Wuhan, where the COVID-19 outbreak was first recorded and very strict lockdown regulations were adopted. At the other five Chinese cities, we also observe a much stronger decrease after the Chinese New Year in 2020 than in 2019. In addition, the duration of the period with low emissions is much longer. Most cities in China display a stronger decrease in 2020 (see Table S1), which is attributed to the COVID-19 measures. The averaged NO_x emission reduction at the selected cities shown in Table S1 is 35%. We also calculate the average reduction of grid cells containing urban areas selected by using the land-use data of the GlobCover Land Cover data set, which was implemented in the CTM by Ding et al. (2015). The inferred emission reduction is about 35% in urban areas, which is the same as the average reduction in the selected cities. Note that the NO_x emissions are usually lower by about 10% during the Chinese New Year with less business and industrial activities (Ding et al., 2017). The timeline of NO_x emissions in Beijing shows a slightly different pattern with a relatively low reduction during the COVID-19 lockdown but already strong emission reductions during the politically important “two-sessions” meeting in March 2019, the most important political meeting of China, and especially the celebration of 70th national anniversary of China around 1 October 2019, when many factories were closed and strict emission regulations were enforced (Yang et al., 2020). Figure 3 also shows that the NO_x emissions start to increase again in March, in line with the step-by-step recovery of the human activities. Except for Wuhan with the emission rebound after 8 April, when the lockdown was lifted, by the end of March all cities reached a level of NO_x emissions close to what was observed in the same period in 2019. This is consistent with the economic target of China that they will accelerate the return to the precrisis economic level after the temporary economic setback due to the COVID-19 outbreak as was reported by Ouyang (2020).




Figure 3. Time series (1 January 2019–28 April 2020) of daily NO_x emissions in seven cities and urban China. Six Chinese cities are considered (Wuhan, Nanjing, Shanghai, Guangzhou, Chongqing, and Beijing) as well as Seoul. The location of Chinese cities is shown in Figure S4.

Besides the urban emissions, we find strong reductions of NO_x emissions from coal power plants. Figure 4 shows time series of NO_x emissions from the Ningxia Province, where the main sources of NO_x are fossil fuel power plants (van der A et al., 2017). Ningxia province can serve as an indication of the national energy production by coal power plants. It has a population of about 6 million, only 0.4% of the total population of China. Its coal production and electricity generation from coal power plants are in the top 10 list of provinces, and about 80% of the generated energy is consumed by the industry (Ningxia Statistics Bureau, 2019). Our inversion results indicate that after the 2020 Chinese New Year, NO_x emissions dropped about 40% in this province, 20% more than in 2019 New Year period. This shows the impact of the COVID-19 regulations on the energy production, especially in the industrial sector. According to the National Bureau of Statistics of China (2020), the total profit of the first 3 months in 2020 made by industrial enterprises decreased around 40% in China compared to the same period of the previous year. The shrinking of the industrial economy results in lower energy consumption, which is clearly reflected by the decrease of NO_x emissions from power plants.




Figure 4. Time series (1 January 2019–28 April 2020) of daily NO_x emissions in Ningxia Province.

4. Surface Concentration Reductions

The effect of the emission reductions on the surface concentration is very relevant for air pollution. In Figure S2 we show the emissions and the modeled surface concentrations from DECSO based on these emissions.




Figure 5. Measured NO_2 concentrations (from 1 January to 12 April 2020) compared to concentrations of the business-as-usual scenario. Cities are chosen similar to Figure 3, except for Seoul. The Chinese New Year is indicated by the blue dashed line.

Although we see a similar time course in both, the reductions in emissions and surface concentrations are different due to the changing meteorology and lifetime of NO_x over time. To further verify the reductions in surface concentrations we used measurements of the in situ stations described in section 2.3. To eliminate the effect of meteorology and transport we compare the measurements of in situ stations with the ensemble model introduced in section 2.4. The model is driven by emission inventories, which are not corrected for the effects of either Spring Festival or the COVID-19 crisis and hence are considered the business-as-usual situation. A possible bias between measurements and model is corrected for by normalizing the results for the first 2 weeks of January. In Figure 5 the ratio between in situ measured NO_2 and the modeled NO_2 is shown. The concentration reductions are shown as green area, while increased concentrations are shown in red. The reduction starts around the Chinese New Year and ends in March. Exception is the concentration level of Wuhan that becomes similar to that of the business-as-usual scenario after the first week of April. Table S1 shows the concentration reduction in P2 compared to P1 for the selected 36 cities. The average concentration reduction is 41%, while for emissions the reduction is 35%. A striking difference between Wuhan and the other Chinese cities is the longer duration (by about 1 month) of the concentration reductions.

5. Conclusions

To study the impact of the COVID-19 regulations on NO_x emissions (one of the key ingredients determining air pollution), we derived daily NO_x emissions at a resolution of $0.25^\circ \times 0.25^\circ$ over East Asia from 2019 to March 2020 by applying the inverse algorithm DECSO to observations from TROPOMI. By grouping the emission into three periods of before, during and after the COVID-19 regulations, we quantified the emission changes on the small spatial scale of city level and from different emission sources such as sea-transport and the energy sector. The observations suggest emission reductions of 20–50% for cities. The emissions reduction of 40% in the Ningxia province reflects the impact of the lockdown measures on the energy sector. Maritime transport is also affected during the COVID-19 regulations, although its emission reductions are dependent on the region. Along the ship track from Shanghai to Guangzhou, the NO_x emissions decreased by 25% during the lockdown and increased again by 18% after the work resumption. While in the region of the Yellow sea and Bohai sea, the emissions decrease by 40% and continued decreasing with another 6% also

in March. To further assess the impact of emission reductions, we compared the in situ NO₂ concentration measurements with simulated surface concentrations from models using unaltered emissions. The emission reductions follow a similar timeline as the surface NO₂ concentrations, which show a sharp reduction around the Chinese New Year and a slow recovery from mid-February to mid-March. Wuhan, the city of the epicenter of the COVID-19 crisis, shows large emission reductions in both February and March, reaching nominal levels in April. In general, we found that activities in the cities returned to normal in March, while as an indicator of the economy, emissions of energy production and international maritime transport took a longer time to return to pre-COVID-19 levels (Table S2).

With the NO_x emissions derived from DECSO using observations from TROPOMI, we are able to get detailed information about the impact on emission changes due to the COVID-19 regulations by accounting for the influence of meteorology, lifetime, and transport of the air pollutants. As the COVID-19 crisis progressively affects all continents, the public health regulations implemented by various countries may have different contributions to air quality. Applying our methodology to different regions can help to quantify the impact of the NO_x emission reductions by the different regulations on not only the improvement of air quality from urban to local to regional scale.

Data Availability Statement

This publication contains modified Copernicus Sentinel-5P data 2019–2020. TROPOMI data are available online (http://www.temis.nl/airpollution/no2col/tropomi_data.php). We acknowledge the ESA GlobCover project for the land use data set (http://due.esrin.esa.int/page_globcover.php). The NO_x emissions data set in this study is available online (http://www.globemission.eu/region_asia/datapage.php?species=NOx_TROPOMI).

Acknowledgments

This research has been supported by the project “Impact study of COVID-19 lockdown measures on air quality and climate” of the European Space Agency (grant number 4000127610/19/I-NS).

References

- Bai, K., Li, K., Guo, J., Yang, Y., & Chang, N.-B. (2020). Filling the gaps of in situ hourly PM_{2.5} concentration data with the aid of empirical orthogonal function analysis constrained by diurnal cycles. *Atmospheric Measurement Techniques*, *13*(3), 1213–1226. <https://doi.org/10.5194/amt-13-1213-2020>
- Bauwens, M., Compernelle, S., Stavrou, T., Müller, J.-F., van Gent, J., Eskes, H., et al. (2020). Impact of coronavirus outbreak on NO₂ pollution assessed using TROPOMI and OMI observations. *Geophysical Research Letters*, *47*, e2020GL087978. <https://doi.org/10.1029/2020GL087978>
- Boersma, K. F., Vinken, G. C. M., & Eskes, H. J. (2016). Representativeness errors in comparing chemistry transport and chemistry climate models with satellite UV–vis tropospheric column retrievals. *Geoscientific Model Development*, *9*(2), 875–898. <https://doi.org/10.5194/gmd-9-875-2016>
- Brasseur, G. P., Xie, Y., Petersen, A. K., Bouarar, I., Flemming, J., Gauss, M., et al. (2019). Ensemble forecasts of air quality in eastern China - Part 1: Model description and implementation of the MarcoPolo-Panda prediction system, version 1. *Geoscientific Model Development*, *12*(1), 33–67. <https://doi.org/10.5194/gmd-12-33-2019>
- Ding, J., Miyazaki, K., Mijling, B., Kurokawa, J. I., Cho, S., Janssens-Maenhout, G., et al. (2017). Intercomparison of NO_x emission inventories over East Asia. *Atmospheric Chemistry and Physics*, *17*(16), 10,125–10,141. <https://doi.org/10.5194/acp-17-10125-2017>
- Ding, J., van der A, R. J., Mijling, B., Jalkanen, J. P., Johansson, L., & Levelt, P. F. (2018). Maritime NO_x emissions over Chinese Seas derived from satellite observations. *Geophysical Research Letters*, *45*, 2031–2037. <https://doi.org/10.1002/2017GL076788>
- Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., & Hao, N. (2015). NO_x emission estimates during the 2014 Youth Olympic Games in Nanjing. *Atmospheric Chemistry and Physics*, *15*(16), 9399–9412. <https://doi.org/10.5194/acp-15-9399-2015>
- Goldberg, D., Lu, Z., Streets, D. G., de Foy, B., Griffin, D., McLinden, C. A., et al. (2019). Enhanced capabilities of TROPOMI NO₂: Estimating NO_x from North American cities and power plants. *Environmental Science & Technology*, *53*(21), 12,594–12,601. <https://doi.org/10.1021/acs.est.9b04488>
- Huang, X., Ding, A., Gao, J., Zheng, B., Zhou, D., Qi, X., et al. (2020). Enhanced secondary pollution offset reduction of primary emissions during COVID-19 lockdown in China. *EarthArXiv*, April 13. <https://doi.org/10.31223/osf.io/hvuzv>
- Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., et al. (2020). Evaluating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island sound. *Atmospheric Measurement Techniques Discussions*, *2020*, 1–52. <https://doi.org/10.5194/amt-2020-151>
- Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., et al. (2020). Abrupt decline in tropospheric nitrogen dioxide over China after the outbreak of COVID-19. *Science Advances*, *6*(28), eabc2992. <https://doi.org/10.1126/sciadv.abc2992>
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., et al. (2013). CHIMERE 2013: A model for regional atmospheric composition modelling. *Geoscientific Model Development*, *6*(4), 981–1028. <https://doi.org/10.5194/gmd-6-981-2013>
- Mijling, B., & van der A, R. J. (2012). Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale. *Journal of Geophysical Research*, *117*, D17302. <https://doi.org/10.1029/2012JD017817>
- Miyazaki, K., Bowman, K., Sekiya, T., Eskes, H., Boersma, F., Worden, H., et al. (2020). An updated tropospheric chemistry reanalysis and emission estimates, TCR-2, for 2005–2018. *Earth System Science Data Discussions*. <https://doi.org/10.5194/essd-2020-30> in review
- Miyazaki, K., Eskes, H. J., & Sudo, K. (2012). Global NO_x emission estimates derived from an assimilation of OMI tropospheric NO₂ columns. *Atmospheric Chemistry and Physics*, *12*(5), 2263–2288. <https://doi.org/10.5194/acp-12-2263-2012>
- National Bureau of Statistics of China (2020). Available online: http://www.stats.gov.cn/english/PressRelease/202004/t20200428_1742015.html (last access date: 3 May 2020)

- NBS, National Bureau of Statistics (2010). *China Statistical Yearbook*. Beijing: China Statistics Press.
- Ningxia Statistics Bureau. (2019). The Achievements of Economic and Social Development during the last 70 years, part 3, in Chinese. Available online: http://tj.nx.gov.cn/tjxx/201909/t20190923_1750612.html (last access date: 9 May 2020)
- Ouyang, S. (2020). COVID-19 will not alter China's growth story, top economic regulator says, *China Daily*. Available online: <https://global.chinadaily.com.cn/a/202004/20/WS5e9d51f8a3105d50a3d1779c.html> (last access date: 27 April 2020)
- Petersen, A. K., Brasseur, G. P., Bouarar, I., Flemming, J., Gauss, M., Jiang, F., et al. (2019). Ensemble forecasts of air quality in eastern China - Part 2: Evaluation of the MarcoPolo-Panda prediction system, version 1. *Geoscientific Model Development*, *12*(3), 1241–1266. <https://doi.org/10.5194/gmd-12-1241-2019>
- Stavrakou, T., Müller, J.-F., Boersma, K. F., van der A, R. J., Kurokawa, J., Ohara, T., & Zhang, Q. (2013). Key chemical NO_x sink uncertainties and how they influence top-down emissions of nitrogen oxides. *Atmospheric Chemistry and Physics*, *13*, 9057–9082. <https://doi.org/10.5194/acp-13-9057-2013>
- Tack, F., Merlaud, A., Iordache, M. D., Pinaridi, G., Dimitropoulou, E., Eskes, H., et al. (2020). Assessment of the TROPOMI tropospheric NO₂ product based on airborne APEX observations. *Atmospheric Measurement Techniques Discussions*, *2020*, 1–55. <https://doi.org/10.5194/amt-2020-148>
- Tian, H., Liu, Y., Li, Y., Wu, C.-H., Chen, B., Kraemer, M. U. G., et al. (2020). An investigation of transmission control measures during the first 50 days of the COVID-19 epidemic in China. *Science*, *368*, 638–642. <https://doi.org/10.1126/science.abb6105>
- van der A, R. J., de Laat, A. T. J., Ding, J., & Eskes, H. J. (2020). Connecting the dots: NO_x emissions along a West Siberian natural gas pipeline. *npj Climate and Atmospheric Science*, *3*(1), 16. <https://doi.org/10.1038/s41612-020-0119-z>
- van der A, R. J., Mijling, B., Ding, J., Koukouli, M. E., Liu, F., Li, Q., et al. (2017). Cleaning up the air: Effectiveness of air quality policy for SO₂ and NO_x emissions in China. *Atmospheric Chemistry and Physics*, *17*(3), 1775–1789. <https://doi.org/10.5194/acp-17-1775-2017>
- van Geffen, J., Boersma, K. F., Eskes, H., Sneep, M., ter Linden, M., Zara, M., & Veefkind, J. P. (2020). S5P TROPOMI NO₂ slant column retrieval: Method, stability, uncertainties and comparisons with OMI. *Atmospheric Measurement Techniques*, *13*(3), 1315–1335. <https://doi.org/10.5194/amt-13-1315-2020>
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., et al. (2012). TROPOMI on the ESA Sentinel-5 precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications. *Remote Sensing of Environment*, *120*, 70–83. <https://doi.org/10.1016/j.rse.2011.09.027>
- Verhoelst, T., Compennolle, S., Pinaridi, G., Lambert, J. C., Eskes, H. J., Eichmann, K. U., et al. (2020). Ground-based validation of the Copernicus Sentinel-5p TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandora global networks. *Atmospheric Measurement Techniques Discussions*, *2020*, 1–40. <https://doi.org/10.5194/amt-2020-119>
- Wang, P., Chen, K., Zhu, S., Wang, P., & Zhang, H. (2020). Severe air pollution events not avoided by reduced anthropogenic activities during COVID-19 outbreak. *Resources, Conservation and Recycling*, *158*, 104814. <https://doi.org/10.1016/j.resconrec.2020.104814>
- WHO. (2020). The World Health Organization. Coronavirus Disease (COVID-19) Pandemic. Available online: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> (last access date: 27 April 2020)
- Yang, Y., Wang, Y., Yao, D., Zhao, S., Yang, S., Ji, D., et al. (2020). Significant decreases in the volatile organic compound concentration, atmospheric oxidation capacity and photochemical reactivity during the National Day holiday over a suburban site in the North China Plain. *Environmental Pollution*, *263*, 114657. <https://doi.org/10.1016/j.envpol.2020.114657>
- Zhang, R., Zhang, Y., Lin, H., Feng, X., Fu, T.-M., & Wang, Y. (2020). NO_x emission reduction and recovery during COVID-19 in East China. *Atmosphere*, *11*(4), 433. <https://doi.org/10.3390/atmos11040433>
- Zhang, Y., & Cao, F. (2015). Fine particulate matter (PM_{2.5}) in China at a city level. *Scientific Reports*, *5*(1), 14884. <https://doi.org/10.1038/srep14884>